

On the Convexity of $\log \det(I + KX^{-1})$

Young-Han Kim and Seung-Jean Kim*

Abstract

A simple proof is given for the convexity of $\log \det(I + KX^{-1})$ in the positive definite matrix variable $X \succ 0$ with a given positive semidefinite $K \succeq 0$.

Convexity of functions of covariance matrices often plays an important role in the analysis of Gaussian channels. For example, suppose $\mathbf{Y}$ and $\mathbf{Z}$ are independent complex Gaussian n -vectors with $\mathbf{Y} \sim N(0, K)$ and $\mathbf{Z} \sim N(0, X)$. Then,

$$I(\mathbf{Y}; \mathbf{Y} + \mathbf{Z}) = \log \det(I + KX^{-1}). \quad (1)$$

The following result is well known in the literature.

Lemma 1. *For a fixed $K \succeq 0$, $\log \det(I + KX^{-1})$ is convex in $X \succ 0$, with strict convexity if $K \succ 0$.*

A simple information theoretic proof was given by Diggavi and Cover [4, Lemma II.3] as a corollary to their main saddle-point theorem stating that the worst additive noise is Gaussian. Very recently Mao *et al.* [6] gave a different, but complicated proof, correcting an incomplete approach taken in Kashyap *et al.* [5].

The main purpose of this note is to introduce an elegant and simple proof technique based on the theory of spectral functions of Hermitian matrices, which will hopefully benefit other problems in information theory with similar structure.

A real-valued function $f(X)$ of Hermitian argument $X \in \mathbb{R}^{n \times n}$ is called a *spectral function* if the value of $f(X)$ depends only on (unordered) eigenvalues of X . If $\lambda(X) \in \mathbb{R}^n$ denotes the ordered eigenvalues of X and g is a real-valued symmetric (=permutation invariant) function on $\mathbb{R}^n$, the composite function $(g \circ \lambda)(X) = g(\lambda(X))$ is a spectral function. Conversely, any spectral function $f(X)$ can be decomposed in this way. It is also easy to see that $f(X)$ is a spectral function if and only if $f(X)$ is unitary invariant, that is, $f(X) = f(QXQ^\dagger)$ for any unitary Q .

Convexity of a spectral function can be checked rather easily; a spectral function $f = g \circ \lambda$ is (strictly) convex if and only if the corresponding symmetric function g is (strictly) convex [3]. In other words, a spectral function $f(X)$ is convex for all Hermitian X if and only if $f(X)$ is convex for all real diagonal X . Examples of convex spectral functions include the trace, the largest eigenvalue, and the sum of the k largest eigenvalues, of a Hermitian matrix; and the trace of the inverse of a positive definite matrix (as well as the log determinant of the

*Email: yhk@ucsd.edu and sjkim@stanford.edu

inverse as will be shown shortly). For a few other interesting properties of spectral functions, refer to [1, Section 5.2].

Now we are ready for a “two-line” proof.

Proof of Lemma 1. Since $X \mapsto K^{1/2}XK^{1/2}$ is a positive linear map (strict if $K \succ 0$) and

$$\log \det(I + KX^{-1}) = \log \det(I + K^{1/2}X^{-1}K^{1/2}),$$

it suffices to establish the strict convexity of

$$f(X) = \log \det(I + X^{-1})$$

in X . (See [2, Section 3.2.4] for the composition rule for convexity.)

Now it is easy to check that $f(X)$ is a spectral function. Indeed,

$$f(X) = g(\lambda(X)) := \sum_{i=1}^n \log \left(1 + \frac{1}{\lambda_i(X)} \right),$$

where $\lambda_1(X), \dots, \lambda_n(X) > 0$ are the eigenvalues of X . But trivially $\log(1 + 1/t)$ is strictly convex for $t > 0$, which implies the strict convexity of g and thus of f . $\square$

References

- [1] J. M. Borwein and A. S. Lewis, *Convex Analysis and Nonlinear Optimization: Theory and Examples*, 2nd ed. New York: Springer, 2006.
- [2] S. Boyd and L. Vandenberghe, *Convex Optimization*. Cambridge: Cambridge University Press, 2004.
- [3] C. Davis, “All convex invariant functions of hermitian matrices,” *Arch. Math.*, vol. 8, pp. 276–278, 1957.
- [4] S. N. Diggavi and T. M. Cover, “The worst additive noise under a covariance constraint,” *IEEE Trans. Inform. Theory*, vol. IT-47, no. 7, pp. 3072–3081, Nov. 2001.
- [5] A. Kashyap, T. Başar, and R. Srikant, “Correlated jamming on MIMO Gaussian fading channels,” *IEEE Trans. Inform. Theory*, vol. IT-50, no. 9, pp. 2119–2123, Sept. 2004.
- [6] W. Mao, X. Su, and X. Xu, “Comments on “Correlated jamming on MIMO Gaussian fading channels”,” *IEEE Trans. Inform. Theory*, vol. IT-52, no. 11, pp. 5163–5165, Nov. 2006.